


## Fundus Diverticulum

### Fundus Divertikülü

A 65-year-old male patient, with no significant medical history, referred to us with melena and fatigue. He used 300 mg acetylsalicylic acid for ischemic heart disease. Gastroscopy was performed as upper gastrointestinal system bleeding was suspected. Widespread erosions in prepyloric area, and an ulcer (Forrest 3), 6-7 mm in diameter, in greater curvature was detected. Erosions were seen in bulb. In addition, the gastroscopy incidentally revealed a diverticulum of 5 mm in diameter in the fundus of stomach (Figure 1,2).

Fundus diverticulum is a rare entity that is mostly asymptomatic but can present with hemorrhage, herniation, reflux, pain or dyspeptic symptoms. Gastroscopy and barium esophagram are generally used for screening. It can be congenital or acquired. For asymptomatic cases, treatment is not required. If symptomatic, treatment is varied according to the severity of symptoms (1).



**Figure 1,2.** Gastrointestinal endoscopic views of fundus diverticulum

Fatih ERMİŞ<sup>1</sup>, Görkem SUCU<sup>2</sup>, Ali KUTLUCAN<sup>2</sup>, Yusuf AYDIN<sup>2</sup>, Sami DOĞAN<sup>3</sup>

<sup>1</sup>Departments of Gastroenterology and <sup>2</sup>Internal Medicine, Düzce University, School of Medicine

<sup>3</sup>Department of General Surgery, Van Research and Educational Hospital, Van

## REFERENCES

1. Lajoie A, Strum WB. Gastric diverticulum presenting as acute hemorrhage. *Gastrointest Endosc* 2008;67:175-6.
2. Simon M, Zuber-Jerger I, Schölmerich J. True gastric diverticulum. *Dig Liver Dis* 2009;41(5):370.
3. Rashid F, Aber A, Iftikhar SY. A review on gastric diverticulum. *World J Emerg Surg* 2012;7:1.

### Correspondence:

Fatih ERMİŞ  
Department of Gastroenterology, Düzce Medical Faculty, Düzce University,  
81620 Konuralp-Düzce • Fax:+90 380 542 13 87  
E-mail: fatihermis2@hotmail.com

Manuscript: 27.11.2013 • Accepted: 01.04.2014